



# Queensland Clinical Guidelines

*Translating evidence into best clinical practice*

## Maternity and Neonatal **Clinical Guideline**

# Trauma in pregnancy

<b>Document title:</b>	Trauma in pregnancy
<b>Publication date:</b>	February 2014
<b>Document number:</b>	MN14.31-V1-R19
<b>Document supplement:</b>	The document supplement is integral to and should be read in conjunction with this guideline.
<b>Amendments:</b>	Full version history is supplied in the document supplement.
<b>Amendment date:</b>	New Document
<b>Replaces document:</b>	New Document
<b>Author:</b>	Queensland Clinical Guidelines
<b>Audience:</b>	Health professionals in Queensland public and private maternity services
<b>Review date:</b>	February 2019
<b>Endorsed by:</b>	Queensland Clinical Guidelines Steering Committee Statewide Maternity and Neonatal Clinical Network (Queensland) Statewide Trauma Network (Queensland)
<b>Contact:</b>	Email: <a href="mailto:MN-Guidelines@health.qld.gov.au">MN-Guidelines@health.qld.gov.au</a> URL: <a href="http://www.health.qld.gov.au/qcg">www.health.qld.gov.au/qcg</a>

#### Disclaimer

These guidelines have been prepared to promote and facilitate standardisation and consistency of practice, using a multidisciplinary approach.

Information in this guideline is current at time of publication.

Queensland Health does not accept liability to any person for loss or damage incurred as a result of reliance upon the material contained in this guideline.

Clinical material offered in this guideline does not replace or remove clinical judgement or the professional care and duty necessary for each specific patient case.

Clinical care carried out in accordance with this guideline should be provided within the context of locally available resources and expertise.

This Guideline does not address all elements of standard practice and assumes that individual clinicians are responsible to:

- Discuss care with consumers in an environment that is culturally appropriate and which enables respectful confidential discussion. This includes the use of interpreter services where necessary
- Advise consumers of their choice and ensure informed consent is obtained
- Provide care within scope of practice, meet all legislative requirements and maintain standards of professional conduct
- Apply standard precautions and additional precautions as necessary, when delivering care
- Document all care in accordance with mandatory and local requirements

© State of Queensland (Queensland Health) 2014



This work is licensed under a Creative Commons Attribution Non-Commercial No Derivatives 3.0 Australia licence. In essence, you are free to copy and communicate the work in its current form for non-commercial purposes, as long as you attribute Queensland Clinical Guidelines, Queensland Health and abide by the licence terms. You may not alter or adapt the work in any way. To view a copy of this licence, visit <http://creativecommons.org/licenses/by-nc-nd/3.0/au/deed.en>

For further information contact Queensland Clinical Guidelines RBWH Post Office, Herston Qld 4029, email [MN-Guidelines@health.qld.gov.au](mailto:MN-Guidelines@health.qld.gov.au), phone (07) 3131 6777. For permissions beyond the scope of this licence contact: Intellectual Property Officer, Queensland Health, GPO Box 48, Brisbane Qld 4001, email [ip\\_officer@health.qld.gov.au](mailto:ip_officer@health.qld.gov.au), phone (07) 3234 1479.

**Flow Chart: Initial assessment and management of the pregnant trauma patient****Principles of care for the pregnant trauma patient**

- Follow ATLS guidelines
- First priority is to treat the woman
- Multidisciplinary team that includes an obstetrician is essential
  - Contact neonatal team early if birth imminent/likely
- Recognise anatomical and physiological changes of pregnancy
- Clear, coordinated and frequent communication essential
- Generally, medications, treatment and procedures as for non-pregnant patient
- Refer pregnant women with major trauma to a trauma centre
  - < 20 weeks gestation: to the nearest trauma centre
  - ≥ 20 weeks gestation: to a trauma centre with obstetric services
- Thoroughly assess all pregnant women – even after minor trauma

**Initial stabilisation**

- As indicated for all trauma patients
- Follow ATLS guidelines
- Initiate early obstetric consultation
- Contact QCC (1300 799 127) to expedite transport & identify receiving facility as required

**Additionally for pregnancy**


- Position (tilt or wedge):
  - Left lateral 15-30° (right side up) *or*
  - Manual displacement of uterus
  - Place wedge under spinal board if necessary
- Routinely administer Oxygen therapy
- Large-bore IV access
- Volume resuscitation (Crystalloid infusion)

**Cardiac arrest**

- Follow ATLS guidelines
- Defibrillate as for non-pregnant patient
- Advanced cardiac life support drugs as indicated for non-pregnant patients
- Perimortem CS if:
  - ≥ 20 weeks gestation
  - No response to effective CPR after 4 minutes

**Abbreviations**

ATLS: Advanced Trauma Life Support  
 CPR: Cardiopulmonary Resuscitation  
 ICS: Caesarean section  
 IETT: Endotracheal tube  
 FAST: Focused Abdominal Sonography for Trauma  
 IV: Intravenous  
 OT: Operating Theatre  
 QCC: Queensland Emergency Medical Coordination Centre  
 >: Greater than  
 ≥: Greater than or equal to



**Flow Chart: Secondary assessment and management of the pregnant trauma patient****Secondary survey****As for non-pregnant patient AND**

- Consult obstetric team
- Maintain high index of suspicion for occult shock and abdominal injury
- Maintain position (tilt or wedge) left lateral 15-30° (right side up) *or*
  - Manual displacement of uterus
  - Wedge spinal board if required
- Obtain obstetric history
  - Gestation
  - Estimated date of delivery
  - Pregnancy complications
- Physical examination
- Assess uterus
  - Tone, rigidity, tenderness
  - Contractions
- Estimate gestational age
  - Fundal height
  - US
  - If uncertain (i.e. severe trauma, no prior US or lack of accurate records) presume viability
- Assess and record FHR
  - Stethoscope *or*
  - Doppler

**Consider - especially for major trauma**

- Rectal examination
- Pelvic exam (obstetric team)
  - Sterile speculum
  - Assess for rupture of membranes, vaginal bleeding, cervical effacement and dilation, cord prolapse, fetal presentation
- Imaging
  - FAST ultrasound
  - Formal obstetric ultrasound
  - Other radiographs
- Blood tests
  - Standard trauma bloods
  - Group and Antibody screen
  - Kleihauer Test if Rh D negative and all women if major trauma (EDTA tube)
  - Consider Coag Profile (major trauma)
- If Rh D negative and ≥ 12 weeks gestation, administer Rh D immunoglobulin (but do not delay definitive care to do so)

**Abbreviations**

CS:	Caesarean section
CTG:	Cardiotocograph
DIC:	Disseminated intravascular coagulopathy
FAST:	Focused Abdominal Sonography for trauma
FHR:	Fetal heart rate
US:	Ultrasound scan
<:	Less than
>:	Greater than
≥:	Greater than or equal to

**Discharge**

- Advise to seek medical advice if:
  - Signs of preterm labour
  - Abdominal pain
  - Vaginal bleeding or discharge
  - Change in fetal movements
- Advise to inform usual maternity care provider of trauma event

Gestation &gt; 24 weeks?

Yes or uncertain

No

Maternal or fetal compromise?

Yes

No

**Consider discharge criteria**

- Obstetric team consulted/agree for discharge
- Reassuring maternal status
- No vaginal loss/bleeding
- Normal CTG/FHR (minimum 4 hours CTG)
  - Interpret CTG with caution at < 28 weeks
- No contractions
- Blood results reviewed
- Rh immunoglobulin given if required
- Social worker referral offered

Discharge criteria met?

Yes

No

**Admit**

- Assess for:
  - Placental abruption
  - Feto-maternal haemorrhage
  - Uterine rupture
  - Preterm labour
  - DIC
- Continuous CTG if > 24 weeks gestation
- Intervene as appropriate
- Consider emergency CS

**Abbreviations**

ATLS	Advanced trauma life support
bpm	Beats per minute
BP	Blood pressure
CPR	Cardiopulmonary resuscitation
CS	Caesarean section
CT	Computerised tomography
CTG	Cardiotocograph
DIC	Disseminated intravascular coagulopathy
ETT	Endotracheal tube
FAST	Focused Abdominal Sonography for Trauma
FHR	Fetal heart rate
FMH	Feto-maternal haemorrhage
FFP	Fresh frozen plasma
INR	International normalised ratio
IV	Intravenous
IVC	Inferior vena cava
mSv	millisievert
MTP	Massive Transfusion Protocol
PPH	Postpartum haemorrhage
QAS	Queensland Ambulance Service
pCO <sub>2</sub>	Partial pressure of carbon dioxide
PT	Prothrombin time
QCC	Queensland Emergency Medical System Coordination Centre
RBWH	Royal Brisbane and Women's Hospital, Brisbane, Queensland
US	Ultrasound scan
rad	Radiation-absorbed dose

**Definitions**

<b>Major trauma</b>	Classification of trauma depends on the mechanism and severity of injury. Refer to Appendix A: Classification of major trauma.
<b>Obstetrician</b>	Local facilities may as required, differentiate the roles and responsibilities assigned in this document to an 'Obstetrician' according to their specific practitioner group requirements; for example to Gynaecologists, General Practitioner Obstetricians, Specialist Obstetricians, Consultants, Senior Registrars and Obstetric Fellows.
<b>Informed consent</b>	When a woman consents to a recommendation about her care after a process of information exchange that involves providing her with sufficient, evidence-based information about all the options for her care so that she can make a decision, in the absence of coercion by any party, that reflects self-determination, autonomy and control. <sup>1</sup>
<b>Sievert</b>	International unit of measurement for the biological effect to human tissue by ionizing radiation.
<b>Woman centred care</b>	<p>Woman centred care includes the affordance of respect and dignity, by supporting the woman to be central and active in her own care<sup>2</sup> through<sup>3</sup>:</p> <ul style="list-style-type: none"> <li>• Holistic care taking account of the woman's physical, psychosocial, cultural, emotional and spiritual needs</li> <li>• Focussing on the woman's expectations, aspirations and needs, rather than the institutional or professional needs</li> <li>• Recognising the woman's right to self determination through choice, control and continuity of care from a known or known caregivers</li> <li>• Recognising the needs of the baby, the woman's family and significant others</li> </ul>

## Table of Contents

1	Introduction .....	7
1.1	Principles of care .....	7
1.2	Patient stratification .....	7
1.3	Family support .....	8
1.4	Transfer and retrieval.....	8
1.5	Clinical standards .....	8
2	Physiological changes in pregnancy .....	9
2.1	Implications for management .....	10
3	Cardiac arrest .....	11
3.1	Perimortem caesarean section .....	11
4	Assessment.....	12
4.1	Primary survey.....	12
4.2	Secondary survey .....	13
4.3	Diagnostic imaging .....	14
5	Obstetric complications .....	15
5.1	Feto-maternal haemorrhage.....	15
5.1.1	Prevention of Rhesus immunisation .....	16
5.2	Preterm labour .....	16
5.3	Placental abruption .....	17
5.4	Uterine rupture.....	18
5.5	Amniotic fluid embolism .....	18
5.6	Disseminated intravascular coagulopathy .....	19
5.7	Musculoskeletal injury.....	19
5.8	Minor trauma.....	20
	References .....	21
	Appendix A: Classification of major trauma in pregnancy .....	24
	Appendix B: Perimortem caesarean section procedure .....	25
	Appendix C: Haemodynamic and laboratory values in pregnancy.....	26
	Appendix D: Seat belt positioning in pregnancy .....	27
	Appendix E: Estimation of gestation.....	28
	Appendix F: Left lateral tilt positioning.....	29
	Appendix G: Approximate fetal effective doses (mSv) from common radiological examinations .....	30
	Acknowledgements.....	31

## List of Tables

Table 1.	Patient category.....	7
Table 2.	Physiological and physical changes in pregnancy .....	9
Table 3.	Implications for management .....	10
Table 4.	Cardiac arrest .....	11
Table 5.	Perimortem caesarean section.....	11
Table 6.	Primary survey additional considerations for pregnancy.....	12
Table 7.	Secondary survey additional considerations for pregnancy.....	13
Table 8.	Diagnostic imaging .....	14
Table 9.	Feto-maternal haemorrhage.....	15
Table 10.	Rh D immunoglobulin .....	16
Table 11.	Preterm labour .....	16
Table 12.	Placental abruption .....	17
Table 13.	Uterine rupture.....	18
Table 14.	Amniotic fluid embolism .....	18
Table 15.	Disseminated intravascular coagulopathy .....	19
Table 16.	Musculoskeletal injury .....	19
Table 17.	Minor trauma.....	20

# 1 Introduction

Trauma affects up to 8% of all pregnancies and is a common cause of non-obstetric maternal morbidity and mortality.<sup>4</sup> Both blunt and penetrating (gunshot or knife related) trauma is encountered in Australia but blunt trauma is the most common. Direct fetal injuries occur in less than 1% of cases of severe blunt abdominal trauma.<sup>5</sup> Even minor injuries in the pregnant woman can be associated with placental abruption, preterm labour, massive feto-maternal haemorrhage, uterine rupture and fetal loss.<sup>4,5</sup> The evidence for care provision is limited with the majority of studies being retrospective and reported outcomes varying widely.<sup>6</sup>

## 1.1 Principles of care

The goal of treatment is maintenance of utero-placental perfusion and fetal oxygenation by avoiding hypoxia and preventing hypotension, acidosis and hypothermia.

- Manage pregnant trauma patients in accordance with the Advanced Trauma Life Support (ATLS) guidelines<sup>6-9</sup>
- The first priority is identification of life threatening injuries to the woman<sup>4,8</sup>
- Thoroughly assess the woman as fetal survival is directly related to maternal wellbeing<sup>4,6</sup>
- A multidisciplinary team approach that includes early involvement of an obstetrician is essential<sup>4,10</sup>
  - Involve neonatal team early if birth imminent/likely
- Recognise maternal anatomical and physiological changes due to pregnancy<sup>4,10</sup>
- Clear, coordinated and frequent communication between care providers is essential<sup>11,12</sup>
- Generally, do not withhold medications, tests, treatments and procedures required for the woman's stabilisation because of pregnancy<sup>6</sup>
- Refer all major trauma cases to a trauma centre [refer to Appendix A: Classification of Major Trauma]
  - If less than 20 weeks gestation, transfer to the nearest trauma centre
  - If greater than or equal to 20 weeks gestation, transfer to a trauma centre with obstetric services<sup>4</sup>
- Provide pregnant women with minor injuries, medical treatment for their injuries and appropriate fetal assessment<sup>13</sup>

## 1.2 Patient stratification

Table 1. Patient category

Category	Considerations
<b>Potentially pregnant</b>	<ul style="list-style-type: none"> <li>• History alone is unreliable in excluding pregnancy</li> <li>• Perform a pregnancy test on all women of child bearing age who experience trauma<sup>5,7,8,14</sup></li> <li>• Where pregnancy is confirmed after a trauma event, provide information and counselling on the implications of the care provided (e.g. diagnostic imaging)</li> </ul>
<b>Pre-viable gestation (&lt; 24 weeks)</b>	<ul style="list-style-type: none"> <li>• Dates and estimations of gestational age may be inaccurate or unreliable</li> <li>• Where there is doubt about the gestation, presume viability</li> <li>• Cardiotocograph (CTG) monitoring not usually indicated</li> <li>• Document presence/absence of fetal heart rate (FHR)</li> </ul>
<b>Viable gestation</b>	<ul style="list-style-type: none"> <li>• Gestations greater than or equal to 24 weeks</li> <li>• Commence CTG monitoring as soon as feasible<sup>15</sup></li> </ul>
<b>Perimortem</b>	<ul style="list-style-type: none"> <li>• Refer to Section 3.1 Perimortem caesarean section (CS)</li> <li>• Refer to Appendix B: Perimortem caesarean section procedure</li> </ul>

### 1.3 Family support

- Share and discuss information with the woman and/or her family in a manner that enables informed choice and consent<sup>16</sup> [refer to Definition of terms and Disclaimer]
- Support a woman centred approach to care and decision making [refer to Definition of terms]
- Provide frequent information about fetal and maternal status to the woman and/or family<sup>11</sup>
  - Explain rationale and risk/benefit for all procedures to enable informed decision making (as circumstances allow)
- Consider intimate partner violence as a cause of trauma in pregnancy<sup>11</sup>
- Offer referral to social workers as appropriate to the circumstances (e.g. intimate partner violence, following fetal demise, if transfer required, for counselling and support)
- Offer debriefing to the woman and/or family following pregnant trauma care events<sup>17</sup>

### 1.4 Transfer and retrieval

- Manage pregnant women at greater than or equal to 20 weeks gestation (or with fundal height higher than umbilicus) who have major trauma, at a Trauma Centre with obstetric services
- In Queensland, Trauma Centres with obstetric services are located at The Townsville Hospital (TTH) and the Royal Brisbane and Women's Hospital (RBWH)
  - If outside the Brisbane greater metropolitan area, arrange inter-hospital transfer via Queensland Emergency Medical System Coordination Centre (QCC)<sup>18</sup>
    - Telephone QCC: 1300 799 127
  - Within the greater metropolitan area of Brisbane, transfer via Queensland Ambulance Service (QAS) to the Royal Brisbane and Women's Hospital (RBWH)
    - Liaise with the RBWH directly – telephone (07) 3646 5900
- Manage pregnant women at less than 20 weeks gestation at a Trauma Centre
  - Arrange transfer/retrieval as per usual local protocols for major trauma
- Where feasible, major trauma surgery should occur in Level 4 or higher operating suite<sup>19</sup>
- Refer to Appendix A: Classification of major trauma in pregnancy

### 1.5 Clinical standards

- Accurate documentation is essential in all cases of maternal collapse, whether or not resuscitation is successful<sup>11,17</sup>
- Consider use of *Queensland Maternity Early Warning Tools* to detect deterioration of pregnant patients<sup>11</sup>
- Review all cases of maternal collapse through the clinical governance process<sup>17</sup>
- Report all maternal deaths as per legislated requirements<sup>20</sup>
- Offer debriefing to clinicians involved in pregnant trauma care events<sup>17</sup>
- Educate clinicians about adaptations to cardiopulmonary resuscitation (CPR) for the pregnant woman<sup>17,21</sup>
- Include information about CPR in the pregnant woman in all generic life support training<sup>17,21</sup>
- Ensure equipment to enable a perimortem CS is accessible in all areas where maternal collapse may occur, including in the Emergency Department<sup>17</sup>
- Provide information to pregnant women about the importance of correct positioning of motor vehicle seat belts while pregnant [refer to Appendix D: Seat belt positioning in pregnancy]



## 2 Physiological changes in pregnancy

An understanding of the anatomic and physiologic alterations of pregnancy is essential.<sup>22</sup> Refer to Appendix C for normal pregnancy values.

Table 2. Physiological and physical changes in pregnancy

	Changes in pregnancy	Implication
<b>Cardiovascular system</b>		
Plasma volume	Increased by up to 50%	Dilutional anaemia Reduced oxygen-carrying capacity Signs of shock due to blood loss appear late
Heart rate	Increased 15–20 bpm	Increased CPR demands
Cardiac output	Increased by 40% Significantly reduced by pressure of gravid uterus on IVC	Increased CPR demands
Uterine blood flow	10% of cardiac output at term	Potential for rapid massive haemorrhage
Systemic vascular resistance	Decreased	Sequesters blood during CPR
Arterial blood pressure (BP)	Decreased by 10–15 mmHg	Decreased reserve
Venous return	Decreased by pressure of gravid uterus on inferior vena cava (IVC)	Increased CPR circulation demands Increased reserve
Coagulation	Increased concentrations of most clotting factors	Activated state of coagulation cascade Increased tendency for thrombosis
<b>Respiratory system</b>		
Respiratory rate	Increased	Decreased buffering capacity, acidosis more likely
Oxygen consumption	Increased by 20%	Hypoxia develops more quickly
Residual capacity	Decreased by 25%	Decreased buffering capacity, acidosis more likely
Arterial pCO <sub>2</sub>	Decreased	Decreased buffering capacity, acidosis more likely
Laryngeal oedema	Increased	Difficult intubation
Mucosal congestion	Increased	Predisposition to airway bleeding
<b>Other changes</b>		
Gastric motility	Decreased	Increased risk of aspiration
Lower oesophageal sphincter	Relaxed	Increased risk of aspiration
Uterus	Enlarged	Diaphragmatic splinting reduces residual capacity and makes ventilation more difficult Aortal compression causes supine hypotension, reduced venous return and significantly impairs CPR Heart rotation to the left – left axis deviation on ECG can be normal in 3 <sup>rd</sup> trimester
Weight	Increased neck and mammary fat levels	Difficult airway management
Pelvic vasculature	Hypertrophied	Potential for massive retroperitoneal haemorrhage with pelvic fracture, uterine trauma
Bowel	Superior displacement	Potential for complex and multiple intestinal injuries with penetrating trauma of the upper abdomen
Bladder	Anterior and superior displacement by uterus	Susceptible to injury as effectively an intra-abdominal organ
Renal blood flow	Increased by 60%. Serum urea, nitrogen, creatinine reduced	'Normal' serum urea nitrogen and creatinine may reflect seriously compromised function

Adapted from Royal College of Obstetricians and Gynaecologists. Maternal collapse in pregnancy and puerperium. Green-top Guideline No. 56. 2011.

## 2.1 Implications for management

Table 3. Implications for management

Aspect	Clinical care
<b>Positioning</b>	<ul style="list-style-type: none"> <li>After 20 weeks gestation, aortocaval compression by the uterus impedes resuscitation by:               <ul style="list-style-type: none"> <li>Decreasing venous return causing supine hypotension</li> <li>Reducing stroke volume and cardiac output<sup>17,23</sup> and</li> <li>Decreasing the effectiveness of thoracic compressions<sup>17,23-25</sup></li> </ul> </li> <li>Position the woman to minimise inferior vena cava (IVC) compression               <ul style="list-style-type: none"> <li>Consider gestation and the ability to provide effective care (e.g. intubation) when determining positioning requirements</li> <li>Left lateral tilt 15–30 degrees<sup>6,8,26,27</sup> (right side up)</li> <li>Place a firm wedge under the right buttock/hip to achieve tilt</li> <li>In cases of major trauma, place the wedge under the spinal board<sup>17</sup></li> </ul> </li> <li>If lateral tilt is not feasible, use manual uterine displacement to minimise IVC compression<sup>5,17,23,26</sup> <ul style="list-style-type: none"> <li>Standing on the woman's left, the clinician places two hands around the uterus and gently pulls the uterus towards themselves<sup>25</sup></li> </ul> </li> <li>Refer to Appendix F: Left lateral tilt positioning</li> </ul>
<b>Common pitfalls</b>	<ul style="list-style-type: none"> <li>Common pitfalls include failure to:               <ul style="list-style-type: none"> <li>Suspect or recognise shock in the presence of normal vital signs</li> <li>Suspect or recognise abdominal injury because of a benign examination</li> <li>Treat shock aggressively with volume replacement (Crystalloids/blood)</li> <li>Suspect and screen for intimate partner violence</li> <li>Recognise and treat supine hypotensive syndrome</li> <li>Conduct necessary radiology studies secondary to fear of injury to the fetus</li> <li>Observe and cardiotocographically monitor all women with minor trauma and a viable fetus (greater than 24 weeks gestation)</li> <li>Detect early pregnancy (by not ordering a urine pregnancy test)</li> <li>Test for Rh D status and administer Rh D immunoglobulin in Rh D negative women</li> <li>Initiate perimortem CS within 4–6 minutes of no response to effective CPR</li> </ul> </li> </ul>

### 3 Cardiac arrest

Table 4. Cardiac arrest

Aspect	Clinical care
<b>Context</b>	<ul style="list-style-type: none"> <li>The efficiency of CPR in maintaining organ perfusion is significantly reduced by aortocaval compression<sup>5,28</sup></li> <li>There is limited evidence about the degree of tilt required to achieve IVC decompression and the effectiveness of chest compressions performed in the left lateral<sup>29</sup></li> </ul>
<b>Management</b>	<ul style="list-style-type: none"> <li>Follow standard guidelines for cardiac arrest<sup>6</sup></li> <li>Position the woman to reduce IVC compression<sup>5</sup> <ul style="list-style-type: none"> <li>Left lateral tilt 15–30 degrees (right side up)</li> <li>Manual displacement of the uterus<sup>29</sup></li> <li>Place wedge under the spinal board if necessary</li> <li>Refer to Section 2.1 Implications for management</li> </ul> </li> <li>Defibrillate as for the non-pregnant trauma patient – no significant shock is delivered to the fetus<sup>25,29</sup> <ul style="list-style-type: none"> <li>Remove CTG leads prior to defibrillation<sup>25,30</sup></li> </ul> </li> <li>Administer advanced cardiac life support drugs as would be indicated for the non-pregnant patient<sup>17,29</sup></li> </ul>

#### 3.1 Perimortem caesarean section

Table 5. Perimortem caesarean section

Aspect	Clinical care
<b>Definition</b>	<ul style="list-style-type: none"> <li>A CS that is initiated after CPR has commenced<sup>31</sup></li> </ul>
<b>Context</b>	<ul style="list-style-type: none"> <li>May improve survival of either or both the woman and fetus<sup>6</sup> but should be considered a resuscitative procedure performed primarily in the interests of maternal survival<sup>17</sup> <ul style="list-style-type: none"> <li>Case studies suggest improved maternal condition/survival results from the increase in venous return after removal of the gravid uterus from the IVC<sup>28,29,31</sup></li> </ul> </li> <li>Survival and neurologic outcome of the viable fetus is related to time between maternal death and birth<sup>14,24,29</sup> <ul style="list-style-type: none"> <li>Best fetal survival occurs when birth is within 4 to 6 minutes of the maternal cardiac arrest<sup>4,29,31</sup></li> <li>Intact fetal survival has not been demonstrated beyond 30 minutes of cardiac arrest<sup>4</sup></li> </ul> </li> <li>Delay in initiating a perimortem CS has been linked to adverse outcomes<sup>11</sup></li> </ul>
<b>Management</b>	<ul style="list-style-type: none"> <li>Where gestation is greater than 20 weeks, perform perimortem CS after 4 minutes of non-response to effective CPR<sup>28</sup></li> <li>Perform CS at the point of resuscitation<sup>17</sup> <ul style="list-style-type: none"> <li>Do not delay perimortem CS by moving the woman to an operating environment or by attempting to assess fetal viability<sup>11,17,28</sup></li> </ul> </li> <li>Continue CPR during and after the procedure<sup>12,28</sup></li> </ul>

## 4 Assessment

Conduct the primary and secondary survey as for non-pregnant patients.<sup>7,10</sup> Additional considerations for pregnancy are outlined in Table 6 and Table 7. Secondary survey

### 4.1 Primary survey

Table 6. Primary survey additional considerations for pregnancy

Aspect	Clinical care
<b>Airway and C-Spine</b>	<ul style="list-style-type: none"> <li>Increased risk of failed intubation – consider:               <ul style="list-style-type: none"> <li>Earlier intubation than for non-pregnant patients<sup>17,22,27,32</sup></li> <li>Use of a short handle laryngoscope</li> <li>Cricoid pressure</li> <li>A smaller endotracheal tube (ETT) due to laryngeal oedema<sup>8,25</sup></li> </ul> </li> <li>Increased risk of aspiration               <ul style="list-style-type: none"> <li>If intubated consider insertion of an orogastric tube<sup>5</sup></li> <li>Consider nasogastric tube if not intubated</li> </ul> </li> <li>Apply cervical spine collar</li> </ul>
<b>Breathing and ventilation</b>	<ul style="list-style-type: none"> <li>Routinely administer supplemental high flow 100% Oxygen<sup>5,8,10,17,27</sup></li> <li>Ventilation volumes may need to be reduced because of elevated diaphragm<sup>25</sup></li> <li>If safe to do so, raise the head of the bed to reduce weight of uterus on the diaphragm and facilitate breathing<sup>32</sup></li> <li>If a chest tube is indicated, place tube 1–2 intercostal spaces above usual fifth intercostal space landmark due to raised diaphragm<sup>5,6,27</sup></li> </ul>
<b>Circulation and haemorrhage control</b>	<ul style="list-style-type: none"> <li>Control obvious external haemorrhage</li> <li>Position with left lateral tilt 15–30 degrees<sup>26,27</sup> (right side up) [refer to Section 2.1]</li> <li>Obtain large-bore intravenous (IV) access               <ul style="list-style-type: none"> <li>Avoid femoral lines due to compression by gravid uterus</li> </ul> </li> <li>Commence Crystalloid IV               <ul style="list-style-type: none"> <li>Assess response – maintain an awareness of pregnancy related physiological parameters</li> <li>Aim to avoid large volumes of crystalloids (greater than 2 L) which may lead to pulmonary oedema due to the relatively low oncotic pressure in pregnancy<sup>12</sup></li> </ul> </li> <li>Avoid vasopressors to restore maternal BP<sup>27</sup> as they may compromise utero-placental flow<sup>33</sup></li> <li>Maintain a high index of suspicion for bleeding and an awareness of the limitations of clinical signs<sup>17</sup></li> <li>Perform a thorough search for occult bleeding as maternal blood flow is maintained at expense of fetus<sup>5</sup></li> <li>Conduct Focused Abdominal Sonography for Trauma (FAST) to assess for intra-abdominal haemorrhage</li> <li>If hypovolaemia is suspected, initiate fluid resuscitation to ensure adequate maternal and utero-placental perfusion<sup>4,17,26</sup></li> <li>Consider Massive Transfusion Protocol (MTP) activation if non-responsive to crystalloids</li> <li>Rapid transfer to operating theatre as indicated</li> <li>Refer to the Queensland Clinical Guideline <i>Postpartum haemorrhage</i> for blood/product replacement and MTP activation protocols<sup>34</sup></li> <li>Evaluate fetal heart rate<sup>26</sup> [refer to Table 7] but do not delay resuscitation for fetal assessments<sup>4</sup></li> </ul>
<b>Disability</b>	<ul style="list-style-type: none"> <li>Rapid neurological evaluation<sup>30</sup> utilising the Glasgow Coma Scale</li> </ul>

## 4.2 Secondary survey

Once the woman is stabilised, further assessment can be undertaken.<sup>6</sup>

Table 7. Secondary survey additional considerations for pregnancy

Aspect	Clinical care
<b>Obstetric history</b>	<ul style="list-style-type: none"> <li>Gestation in weeks/estimated date of delivery</li> <li>Previous pregnancy complications</li> <li>Prenatal care</li> <li>History of vaginal bleeding</li> </ul>
<b>Physical Examination</b>	<ul style="list-style-type: none"> <li>Head to toe examination as for non-pregnant trauma patients<sup>26</sup></li> <li>Inspect abdomen for ecchymosis or asymmetry</li> <li>In cases of motor vehicle accident, incorrect positioning of the seat belt across the gravid uterus may [refer to Appendix D: Seat belt positioning in pregnancy]:             <ul style="list-style-type: none"> <li>Cause marked bruising of the abdomen</li> <li>Increase the risk of placental abruption</li> <li>Increase the risk of uterine rupture</li> </ul> </li> <li>Assess uterine tone, contractions, rigidity, tenderness, palpable fetal parts             <ul style="list-style-type: none"> <li>The gravid abdomen may be relatively insensate to peritoneal irritation</li> </ul> </li> </ul>
<b>Estimation of gestational age</b>	<ul style="list-style-type: none"> <li>Can be estimated by measuring fundal height             <ul style="list-style-type: none"> <li>Measure the vertical distance in the midline from the symphysis pubis to the top of the fundus in centimetres. This measurement correlates approximately with the gestational age</li> <li>Refer to Appendix E: Estimation of gestational age</li> </ul> </li> <li>Ultrasound scan (US) estimation<sup>31</sup> <ul style="list-style-type: none"> <li>Biparietal diameter (BPD) of 60 mm generally corresponds to a gestation age of approximately 24 weeks</li> </ul> </li> <li>Mark the top of the fundus to evaluate the possibility of concealed abruption as noted by increasing fundal height<sup>10</sup></li> </ul>
<b>Fetal heart rate monitoring</b>	<ul style="list-style-type: none"> <li>Normal FHR 110–160 bpm<sup>35</sup></li> <li>FHR can be assessed using standard stethoscope from about 20 weeks and Doppler from about 12 weeks<sup>5,33</sup> <ul style="list-style-type: none"> <li>Differentiate maternal and FHR as maternal tachycardia may cause confusion<sup>26</sup></li> </ul> </li> <li>For gestations greater than 24 weeks (major trauma), initiate continuous cardiotocography (CTG) as soon as feasible<sup>5,26</sup> <ul style="list-style-type: none"> <li>Good sensitivity for immediate adverse outcome</li> <li>Detects uterine irritability and abnormal fetal heart rate patterns</li> </ul> </li> <li>Abnormalities may be the only indication of injury or compromise to the fetus<sup>27</sup> <ul style="list-style-type: none"> <li>Persistent fetal bradycardia more than 5 minutes, loss of baseline variability or recurrent complex variable or late decelerations indicates fetal compromise<sup>35</sup></li> <li>Sinusoidal trace indicates fetal anaemia</li> </ul> </li> <li>CTG application and interpretation requires clinicians trained in their use             <ul style="list-style-type: none"> <li>Physiological control of FHR and resultant CTG trace interpretation differs in the preterm fetus compared to the term fetus, especially at gestations less than 28 weeks<sup>36</sup></li> <li>CTG trace review should be performed by a clinician experienced and confident with CTG interpretation relevant to the gestation<sup>36</sup></li> <li>Move staff and equipment to the woman's location rather than transporting a woman to an obstetric unit for monitoring</li> </ul> </li> </ul>
<b>Pelvic/vaginal examination</b>	<ul style="list-style-type: none"> <li>If major trauma, perform a rectal examination to assess for spinal cord damage or local trauma</li> <li>Perform sterile speculum vaginal examination<sup>8,26</sup> as clinically indicated (preferably by an obstetric/maternity team member<sup>8,26</sup>)             <ul style="list-style-type: none"> <li>Evaluate for ruptured membranes, vaginal bleeding, cord prolapse, cervical effacement and dilation in labour, fetal presentation<sup>8</sup></li> <li>Vaginal bleeding may indicate preterm labour, abruption, pelvic fracture or uterine rupture<sup>10</sup></li> </ul> </li> <li>Consider urinary catheter insertion<sup>32</sup></li> </ul>

### 4.3 Diagnostic imaging

Table 8. Diagnostic imaging

Aspect	Clinical care
<b>Context</b>	<ul style="list-style-type: none"> <li>The fetus is most vulnerable to radiation during the first 15 weeks of gestation<sup>37</sup></li> <li>The risks of radiation to the fetus are small compared with the risk of missed or delayed diagnosis of trauma<sup>38</sup></li> <li>Increased risks to the embryo or fetus have not been observed for intellectual disability, birth defects, growth restriction, neurobehavioural effects, impaired school performance, convulsive disorders, or embryonic or fetal death below an effective dose of 100 mSv<sup>39</sup></li> <li>Although iodinated contrast agents cross the placenta and may be taken up by the fetal thyroid, no cases of fetal goitre or abnormal neonatal thyroid function have been reported in connection with in-utero contrast exposure<sup>4</sup></li> <li>Gadolinium has known teratogenic effects on animals and is not recommended unless benefits clearly outweigh the risks<sup>40</sup></li> </ul>
<b>Management</b>	<ul style="list-style-type: none"> <li>X-ray examinations of the extremities, head and skull, mammography and computerised tomography (CT) examinations of the head and neck can be undertaken on pregnant or possibly pregnant women without concern<sup>39,41</sup></li> <li>Other X-ray examinations may also be undertaken if the radiation dose to the embryo or fetus is likely to be less than 1 mSv<sup>41,42</sup></li> <li>Where a procedure on a pregnant woman may result in a radiation dose of more than 1 mSv to an embryo or fetus, the following is required<sup>41,42</sup>: <ul style="list-style-type: none"> <li>Be justified on an individual basis</li> <li>Include an assessment of the risks to the: <ul style="list-style-type: none"> <li>Embryo or fetus from radiation exposure</li> <li>Woman if the procedure is not performed</li> </ul> </li> <li>An estimate of the expected radiation dose to the embryo or fetus is made and documented in the health record <ul style="list-style-type: none"> <li>If practicable, consult a medical physicist if individual estimation/calculation of embryo or fetal dose is required</li> </ul> </li> </ul> </li> <li>Optimisation of the examination's exposure parameters has the largest effect on doses</li> <li>Personal protective equipment, (e.g. lead gown) is advised for pregnant women only when the position of the uterus is in the direct X-ray beam (and not if it interferes with imaging)<sup>41</sup></li> <li>It is preferable to perform a single CT scan with iodinated contrast rather than perform multiple suboptimal studies without contrast<sup>4</sup></li> <li>Refer to Appendix G: Approximate fetal effective doses (mSv) arising from common radiological examination of pregnant women</li> <li>Provide information and counselling to women exposed to radiation during diagnosis and care<sup>43</sup></li> <li>Refer to local Radiation Safety and Protection Plans</li> </ul>
<b>Ultrasound</b>	<ul style="list-style-type: none"> <li>US can assess solid organ injury, intra-peritoneal fluid, gestational age, FHR, fetal activity, fetal presentation, placental location, amniotic fluid volume<sup>4,38</sup></li> <li>US is not a reliable indicator of recent placental abruption<sup>4,38,44</sup></li> <li>FAST scan is as accurate as in non-pregnant patients<sup>8</sup> for intra-abdominal free fluid</li> <li>Consider formal obstetric US following FAST as clinically indicated</li> </ul>

## 5 Obstetric complications

### 5.1 Feto-maternal haemorrhage

Table 9. Feto-maternal haemorrhage

Aspect	Clinical care
<b>Context</b>	<ul style="list-style-type: none"> <li>Feto-maternal haemorrhage (FMH) occurs in approximately 10–30% of pregnant trauma patients<sup>5,30</sup></li> <li>The severity of the FMH is related to the size of the bleed in relation to the overall fetal blood volume, the rate at which this blood is lost and whether the event is acute or chronic</li> <li>Clinical presentation of FMH is variable and can be non-specific<sup>45,46</sup> <ul style="list-style-type: none"> <li>Decreased or absent fetal movements have been reported<sup>45,46</sup></li> <li>Fetal distress – especially if the fetal heart tracing is sinusoidal (indicating fetal anaemia)</li> <li>Massive FMH is a rare but severe complication which can result in fetal anaemia, fetal hypoxia, intrauterine death or neonatal neurologic damage<sup>46</sup></li> <li>Women may experience a transfusion reaction (nausea, oedema, fever, and chills)<sup>45</sup></li> <li>May occur more commonly with anteriorly located placentae and in women who experience uterine tenderness after trauma<sup>47</sup></li> </ul> </li> </ul>
<b>Assessment of feto-maternal haemorrhage</b>	<ul style="list-style-type: none"> <li>The Kleihauer test is used to detect and quantify FMH<sup>48</sup> <ul style="list-style-type: none"> <li>Commonly to determine dose of Rh D immunoglobulin for Rh D negative women<sup>49</sup></li> <li>Results are reported quantitatively in mL of fetal blood within maternal circulation</li> <li>A 'negative' result is commonly understood to be less than 1 mL of fetal blood</li> <li>The Kleihauer test is not a test for placental abruption<sup>44,50</sup></li> <li>The evidence is limited about the usefulness of a positive Kleihauer test for predicting outcomes and guiding clinical management<sup>49,51-53</sup> (beyond determining the dose of Rh D immunoglobulin for Rh D negative women)</li> </ul> </li> <li>Flow cytometry is the most accurate quantitative test for FMH<sup>48</sup> and will be initiated by Pathology Queensland as a standard procedure when the quantitative result of the Kleihauer test is greater than 4 mL</li> </ul>
<b>Management</b>	<ul style="list-style-type: none"> <li>Continuous electronic fetal monitoring of the viable fetus</li> <li>Abdominal US to detect fetal heart activity, placental location, amniotic fluid index, suspected intraperitoneal bleeding, gestational age, fetal weight</li> <li>Elevated peak systolic velocity of the fetal middle cerebral artery correlates with fetal anaemia<sup>54,55</sup></li> <li>Emergency CS may be indicated</li> </ul>
<b>Recommendation</b>	<ul style="list-style-type: none"> <li>Following a trauma event: <ul style="list-style-type: none"> <li>Kleihauer test is recommended for all Rh D negative women greater than 12 weeks gestation<sup>48,56,57</sup> to determine the dose of Rh D immunoglobulin required [refer to Table 10]</li> <li>Consider a Kleihauer test for all women with major or abdominal trauma to aid identification of FMH and inform immediate and longer term pregnancy management and outcomes</li> <li>Maintain a high index of suspicion and clinical surveillance for the possibility of significant FMH</li> </ul> </li> </ul>

### 5.1.1 Prevention of Rhesus immunisation

Table 10. Rh D immunoglobulin

Aspect	Clinical care
<b>Assessment</b>	<ul style="list-style-type: none"> <li>For the Rh D negative woman greater than 12 weeks gestation, collect maternal blood (blood group, antibody screen and Kleihauer test) prior to administration of Rh D immunoglobulin<sup>56</sup></li> <li>Do not delay or withhold administration of Rh D immunoglobulin based on or pending the results of quantitative testing</li> </ul>
<b>Rh D immunoglobulin</b>	<ul style="list-style-type: none"> <li>Indicated for the non-sensitised Rh D negative woman within 72 hours of the sensitising event where:               <ul style="list-style-type: none"> <li>Gestation is greater than 12 weeks</li> <li>Gestation is unknown/possibly greater than 12 weeks</li> </ul> </li> <li>Not indicated when gestation is less than 12 weeks</li> <li>If not offered within 72 hours, a dose offered within 9–10 days may provide protection<sup>56</sup></li> <li>625 IU of Rh D immunoglobulin protects against 6 mL fetal red cells (12 mL whole blood), which is equivalent to 0.25% fetal cells in the maternal circulation<sup>48</sup></li> </ul>
<b>Dose</b>	<ul style="list-style-type: none"> <li>Rh D immunoglobulin 625 IU via intramuscular injection<sup>48,56</sup></li> <li>If FMH is quantified at greater than 6 mL, give additional doses of Rh D immunoglobulin sufficient to provide immunoprophylaxis within 72 hours<sup>56</sup> (625 IU for each additional 6 mL (or part thereof) of fetal red cells detected)</li> </ul>
<b>Contraindications</b>	<ul style="list-style-type: none"> <li>Rh D positive woman</li> <li>Rh D negative woman with preformed Anti-D antibodies<sup>56</sup></li> <li>Previous sensitivity or allergy to Rh D immunoglobulin</li> </ul>

## 5.2 Preterm labour

Table 11. Preterm labour

Aspect	Clinical care
<b>Context</b>	<ul style="list-style-type: none"> <li>Onset of labour before 37 completed weeks gestation<sup>50</sup></li> </ul>
<b>Clinical presentation</b>	<ul style="list-style-type: none"> <li>Uterine contractions of more than 4 per hour accompanied by cervical change<sup>8</sup></li> <li>Cramping abdominal/back pain<sup>50</sup></li> <li>Pelvic pressure<sup>50</sup></li> <li>An increase or change in vaginal discharge<sup>50</sup></li> <li>Vaginal bleeding<sup>50</sup></li> </ul>
<b>Management</b>	<ul style="list-style-type: none"> <li>Consult with an obstetrician regarding management appropriate for the circumstances</li> <li>Refer to the Queensland Clinical Guideline <i>Preterm Labour</i><sup>58</sup>:               <ul style="list-style-type: none"> <li>Consider tocolytic therapy</li> <li>Consider corticosteroids aimed at promoting fetal lung maturity</li> </ul> </li> </ul>



### 5.3 Placental abruption

Table 12. Placental abruption

Aspect	Clinical care
<b>Context</b>	<ul style="list-style-type: none"> <li>Common complication of trauma especially following motor vehicle accidents<sup>15</sup> (rate in general obstetrical population of 0.4 to 1.3%)<sup>59</sup> <ul style="list-style-type: none"> <li>One study reported frequency after motor vehicle accident with severe, non-severe or no injury of 13%, 7.4% and 8.5% respectively<sup>59</sup></li> </ul> </li> <li>Leading cause of fetal death following trauma<sup>4,15</sup> accounting for 50–70% of all trauma-related fetal losses<sup>4</sup></li> <li>Can occur with rapid deceleration without direct trauma<sup>7</sup></li> <li>Can occur following relatively minor trauma<sup>7,47</sup></li> <li>Has not been reported when less than 1 contraction is present in any 10 minute interval over a 4 hour period<sup>6</sup></li> </ul>
<b>Clinical presentation</b>	<ul style="list-style-type: none"> <li>Abdominal pain<sup>44,60</sup></li> <li>Vaginal bleeding<sup>60</sup> – 80% of cases<sup>50</sup> <ul style="list-style-type: none"> <li>Amount does not necessarily correlate with severity<sup>50</sup></li> </ul> </li> <li>Uterine contractions<sup>4</sup></li> <li>Uterine tenderness<sup>60</sup>/tense or 'woody' feel<sup>44</sup></li> <li>Expanding fundal height<sup>10</sup></li> <li>Evidence of fetal compromise<sup>44</sup></li> <li>Maternal haemodynamic instability<sup>50</sup></li> <li>Can also present asymptotically<sup>50</sup></li> </ul>
<b>Investigations</b>	<ul style="list-style-type: none"> <li>Although US may detect abruption, it is not sensitive enough to exclude abruption<sup>44,60</sup> <ul style="list-style-type: none"> <li>False negative reported 50–80%<sup>38</sup></li> </ul> </li> <li>CTG is better than US in risk stratifying for suspected placental abruption<sup>4,5</sup> <ul style="list-style-type: none"> <li>Uterine contractions have high-frequency, low-amplitude pattern with an elevated baseline tone<sup>50</sup></li> <li>Fetal heart rates can show recurrent late or variable decelerations, bradycardia, or sinusoidal patterns<sup>50</sup></li> </ul> </li> <li>Consider feto-maternal haemorrhage<sup>17</sup> [refer to Table 10]</li> <li>Request full blood count, coagulation studies, blood group and antibody<sup>50</sup></li> </ul>
<b>Management</b>	<ul style="list-style-type: none"> <li>Difficult to diagnose in mild forms<sup>12,15</sup></li> <li>Consider admission for surveillance as clinically indicated</li> <li>Give Rh D immunoglobulin to all non-sensitised Rh D negative women independent of whether routine antenatal prophylactic Rh D immunoglobulin has been administered<sup>17</sup> [refer to Table 9]</li> <li>Consider antenatal corticosteroids between 24 and 34 weeks + 6 days gestation<sup>17</sup></li> <li>Monitor for disseminated intravascular coagulopathy (DIC) and request urgent clotting studies, platelet count as indicated<sup>44</sup> <ul style="list-style-type: none"> <li>Do not delay treatment by waiting for coagulation results if massive blood loss occurs<sup>44</sup></li> </ul> </li> <li>Significant placental abruption requires urgent delivery by CS<sup>44</sup> <ul style="list-style-type: none"> <li>Incision – mid line preferable if other abdominal injuries suspected</li> <li>Refer to Queensland Clinical Guideline <i>Postpartum haemorrhage</i> for management of PPH, blood/product replacement and MTP activation protocols<sup>34</sup></li> </ul> </li> </ul>

## 5.4 Uterine rupture

Table 13. Uterine rupture

Aspect	Clinical care
<b>Context</b>	<ul style="list-style-type: none"> <li>Uterine rupture is more likely with advanced gestational age and severe direct abdominal trauma<sup>5</sup></li> <li>Diagnosis usually made on US<sup>50</sup> (extrusion of uterine contents, free fluid in pelvis)</li> </ul>
<b>Clinical presentation</b>	<ul style="list-style-type: none"> <li>CTG abnormalities<sup>12,50</sup> (most common feature)</li> <li>Fetal demise<sup>50</sup></li> <li>Positive FAST</li> <li>Uterine tenderness/pain<sup>50</sup></li> <li>Vaginal bleeding<sup>50</sup></li> <li>Palpable fetal parts<sup>50</sup></li> <li>Maternal shock including hypotension and tachycardia<sup>50</sup></li> </ul>
<b>Management</b>	<ul style="list-style-type: none"> <li>CS with midline laparotomy</li> <li>Urgent delivery of fetus</li> <li>Repair of uterus (simple repair, subtotal hysterectomy or total hysterectomy) as indicated by individual circumstances<sup>12</sup></li> <li>Prompt haemodynamic resuscitation with blood products decreases risk of DIC<sup>61</sup></li> <li>Hysterectomy if uncontrolled haemorrhage<sup>61</sup></li> </ul>

## 5.5 Amniotic fluid embolism

Table 14. Amniotic fluid embolism

Aspect	Clinical care
<b>Context</b>	<ul style="list-style-type: none"> <li>Exposure of the amniotic fluid to the maternal circulation may cause amniotic fluid embolism and DIC<sup>5</sup> although the exact mechanism is unknown<sup>50</sup></li> </ul>
<b>Clinical presentation</b>	<ul style="list-style-type: none"> <li>Maternal hypotension<sup>17,50</sup> (100% of women<sup>12,25</sup>)</li> <li>Respiratory distress</li> <li>Seizure<sup>17</sup></li> <li>Cardiac arrest<sup>17,50</sup> (87% of women<sup>25</sup>)</li> <li>Fetal distress develops acutely<sup>17,50</sup></li> <li>Massive haemorrhage<sup>17</sup></li> <li>Coagulopathy/DIC<sup>17,50</sup></li> </ul>
<b>Management</b>	<ul style="list-style-type: none"> <li>Supportive care – there is no proven effective treatment<sup>12,17,25</sup></li> <li>Resuscitation and airway management<sup>50</sup></li> <li>Multidisciplinary care</li> <li>Blood product replacement including Fresh Frozen Plasma (FFP), Platelets and Cryoprecipitate<sup>50</sup></li> </ul>

## 5.6 Disseminated intravascular coagulopathy

Table 15. Disseminated intravascular coagulopathy

Aspect	Clinical care
<b>Context</b>	<ul style="list-style-type: none"> <li>May arise following placental abruption, fetal demise and amniotic fluid embolism<sup>12</sup></li> <li>Early delivery protects against severe DIC – which is partly due to the massive release of thromboplastins from the damaged uterus<sup>12</sup></li> </ul>
<b>Clinical presentation</b>	<ul style="list-style-type: none"> <li>May result in clinically detectable microvascular bleeding as well as abnormal blood coagulation tests<sup>34,62</sup> including:               <ul style="list-style-type: none"> <li>Platelet count less than <math>50 \times 10^9/L</math></li> <li>Prothrombin time (PT) greater than 1.5 x normal</li> <li>International normalised ratio (INR) greater than 1.5</li> <li>Activated partial thromboplastin time (aPTT) greater than 1.5 x normal</li> <li>Fibrinogen level less than 2.5 g/L<sup>34</sup></li> </ul> </li> </ul>
<b>Management</b>	<ul style="list-style-type: none"> <li>Refer to Queensland Clinical Guideline <i>Primary postpartum haemorrhage</i> for management, blood/product replacement and MTP activation protocols<sup>34</sup></li> <li>Treat underlying cause</li> <li>Requires early aggressive management<sup>17</sup></li> <li>Collect baseline bloods early and frequently</li> <li>If clinical signs present do not delay treatment by waiting for coagulation results<sup>44</sup></li> <li>Avoid hypothermia and acidosis</li> <li>If undelivered, deliver fetus and placenta<sup>17</sup></li> <li>Advise Platelet transfusion if marked or moderate thrombocytopenia</li> <li>Advise early use of Cryoprecipitate to maintain fibrinogen levels above 2.5 g/L</li> <li>Give FFP if actively bleeding or significantly elevated INR</li> <li>Consult with a Haematologist<sup>44</sup>, especially if considering:               <ul style="list-style-type: none"> <li>Recombinant Activated Factor VII (rFVIIa) – has been used off licence in some obstetric patients with DIC</li> <li>Tranexamic Acid</li> </ul> </li> </ul>

## 5.7 Musculoskeletal injury

Management principles are generally the same as for the non-pregnant patient.

Table 16. Musculoskeletal injury

Type	Clinical care
<b>Penetrating trauma</b>	<ul style="list-style-type: none"> <li>Low threshold for exploratory laparotomy<sup>10</sup></li> </ul>
<b>Spine and spinal cord injuries</b>	<ul style="list-style-type: none"> <li>Adequate immobilisation of neck and spine<sup>10</sup></li> <li>Position left lateral tilt 15–30° (right side up) – if possible</li> <li>Early multidisciplinary approach to care</li> <li>Consider delivery at advanced gestations</li> </ul>
<b>Major pelvic fracture</b>	<ul style="list-style-type: none"> <li>Immobilise pelvis</li> <li>Vaginal birth is not absolutely contraindicated<sup>10</sup> <ul style="list-style-type: none"> <li>Birth by CS if unstable fracture or pelvic architecture disrupted<sup>10</sup></li> </ul> </li> <li>Consider fetal injury/skull fracture – may be more common with fetal head engagement<sup>10</sup> <ul style="list-style-type: none"> <li>Consult with neonatologist</li> </ul> </li> </ul>
<b>Limb fracture and longer term immobility</b>	<ul style="list-style-type: none"> <li>Assess for venous thromboembolism (VTE) risk and consider prophylaxis<sup>37</sup> <ul style="list-style-type: none"> <li>Refer to the Queensland Clinical Guideline <i>Venous thromboembolism (VTE) prophylaxis in pregnancy and the puerperium</i><sup>63</sup></li> </ul> </li> </ul>

## 5.8 Minor trauma

Table 17. Minor trauma

Aspect	Clinical care
<b>Definition</b>	<ul style="list-style-type: none"> <li>Any trauma injury that does not meet the criteria for defining major trauma</li> <li>Refer Appendix A Classification of major trauma in pregnancy</li> </ul>
<b>Context</b>	<ul style="list-style-type: none"> <li>Severity of injury may not be predictive of fetal outcome<sup>4,13</sup></li> <li>Adverse fetal outcomes are increased after minor trauma not requiring hospitalisation<sup>4,7,13</sup></li> <li>Placental abruption has not been reported when less than one contraction is present in any 10 minute interval over a 4 hour period<sup>6</sup></li> </ul>
<b>FHR monitoring</b>	<ul style="list-style-type: none"> <li>CTG provides good screening/high sensitivity for <i>immediate</i> adverse outcome</li> <li>Monitor FHR via CTG for 4 hours<sup>26,38,61</sup> at a minimum</li> </ul>
<b>Discharge following minor trauma</b>	<ul style="list-style-type: none"> <li>Consult with the obstetric team prior to discharge</li> <li>Criteria:             <ul style="list-style-type: none"> <li>Normal CTG<sup>26</sup> <ul style="list-style-type: none"> <li>Interpret with caution at 24–28 weeks gestation</li> <li>Refer to Table 7 for Fetal heart rate monitoring considerations</li> </ul> </li> <li>No contractions</li> <li>No vaginal bleeding/loss<sup>26</sup></li> <li>Reassuring maternal status</li> <li>Laboratory evaluation within normal limits</li> <li>Kleihauer test reviewed and sufficient Rh D immunoglobulin administered (if required)</li> </ul> </li> <li>Offer social work referral before discharge</li> <li>Advise the woman to inform her usual obstetric care provider of the trauma event</li> <li>Increased antenatal surveillance is required even after minor trauma as the risk of adverse obstetric outcomes is increased including premature labour, low birth weight, fetal demise<sup>13</sup> and placental abruption<sup>4</sup></li> <li>Advise the woman to inform her usual obstetric care provider of the trauma event</li> <li>Advise the woman to seek medical advice if experiencing:             <ul style="list-style-type: none"> <li>Signs of preterm labour</li> <li>Abdominal pain</li> <li>Vaginal bleeding</li> <li>Change in fetal movements</li> </ul> </li> </ul>

## References

1. National Health and Medical Research Council (NHMRC), Department of Health and Ageing, Australian Government. National guidance on collaborative maternity care. Canberra: NHMRC; 2010.
2. NHS, Quality Improvement Scotland. Pathways for maternity care. Keeping Childbirth Natural and Dynamic Programme. 2009.
3. Homer C, Brodie P, Leap N. Midwifery continuity of care: a practical guide. Sydney: Elsevier; 2008.
4. Brown S, Mozurkewich E. Trauma during pregnancy. *Obstetrics and Gynecology Clinics of North America*. 2013; 40(1):47.
5. McAuley DJ. Trauma in pregnancy: anatomical and physiological considerations. *Trauma*. 2004; 6(4):293-300.
6. Mendez-Figueroa H, Dahlke JD, Vrees RA, Rouse DJ. Trauma in pregnancy: an updated systematic review. *Am J Obstet Gynecol*. 2013; vol 209(1):1-10.
7. Sela HY, Weiniger CF, Hersch M, Smueloff A, Laufer N, Einav S. The pregnant motor vehicle accident casualty: adherence to basic workup and admission guidelines. *Annals of Surgery*. 2011; 254(2):346-352.
8. Einav S, Sela HY, Weiniger CF. Management and outcomes of trauma during pregnancy. *Anesthesiol Clin*. 2013; 31(1):141-56.
9. American College of Surgeons Committee on Trauma. Advanced Trauma Life Support for Doctors, Student Manual. Chicago: First Impressions; 2008.
10. Muench M, Canterino J. Trauma in pregnancy. *Obstetrics and Gynecology Clinics of North America*. 2007; 34:555-583.
11. Centre for Maternal and Child Enquiries (CMACE). Saving mothers' lives: reviewing maternal deaths to make motherhood safer. 2006-2008. The eighth report on confidential enquiries into maternal deaths in the United Kingdom. *BJOG*. 2011; 118 (Suppl. 1):1-203.
12. Grady K, Howell C, Cox C, editors. The MOET course manual: managing obstetric emergencies and trauma. 2nd ed. London: RCOG press; 2007.
13. Fischer PE, Zarzaur BL, Fabian TC, Magnotti LJ, Croce MA. Minor trauma is an unrecognized contributor to poor fetal outcomes: a population-based study of 78,552 pregnancies. *The Journal of Trauma*. 2011; 71(1):90-93.
14. Barraco RD, Simon BJ, Weiss PM, Chiu WC, Clancy TV, Como JJ, et al. Practice management guidelines for the diagnosis and management of injury in the pregnant patient: the EAST practice management guidelines work group. *The Journal of Trauma*. 2010; 69(1):211-214.
15. Wyant AR, Collett D. Trauma in pregnancy: diagnosis and management of two patients in one. *JAAPA*. 2013; 26(5):24-9.
16. Queensland Health. Guide to informed decision-making in healthcare. 2012 [cited 2013 October 08]. Available from: <http://www.health.qld.gov.au/consent/documents/ic-guide.pdf>.
17. Royal College of Obstetricians and Gynaecologists. Maternal collapse in pregnancy and puerperium. Green-top Guideline No. 56. 2011.
18. Queensland Government, Statewide Clinical Coordination and Retrieval Services. SOP No.3.7 Criteria for early notification of trauma for interfaculty transfers. 2010 [cited 2013 October 2]. Available from: [http://qheps.health.qld.gov.au/ccrs/sops/sop\\_index.htm](http://qheps.health.qld.gov.au/ccrs/sops/sop_index.htm).
19. Queensland Government, Clinical Access and Redesign Unit. Perioperative services. In: Clinical services capability framework for public and licensed private health facilities v3.1: Queensland Health; 2012.
20. Queensland Government. Public Health Act 2005. [cited 2013 December 01]. Available from: <https://www.legislation.qld.gov.au/LEGISLTN/CURRENT/P/PubHealA05.pdf>.
21. Smith A, Edwards S, Siassakos D. Effective team training to improve outcomes in maternal collapse and perimortem caesarean section. *Resuscitation*. 2012; 83(10):1183-1184.

22. Suresh MS, Latoya Mason C, Munnur U. Cardiopulmonary resuscitation and the parturient. *Best Practice and Research: Clinical Obstetrics and Gynaecology*. 2010; 24(3):383-400.
23. Kim S, You JS, Lee HS, Lee JH, Park YS, Chung SP, et al. Quality of chest compressions performed by inexperienced rescuers in simulated cardiac arrest associated with pregnancy. *Resuscitation*. 2013; 84(1):98-102.
24. Guven S, Yazar A, Yakut K, Aydogan H, Erguve M, Avci E. Postmortem cesarean: report of our successful neonatal outcomes after severe trauma during pregnancy and review of the literature. *The Journal of Maternal-Fetal Medicine*. 2012; 25(7):1102-04.
25. Jones R, Baird SM, Thurman S, Gaskin IM. Maternal cardiac arrest: an overview. *Journal of Perinatal and Neonatal Nursing*. 2012; 26(2):117-123.
26. Chames MC, Pearlman MD. Trauma during pregnancy: outcomes and clinical management. *Clinical Obstetrics & Gynecology*. 2008; 51(2):398-408.
27. Meroz Y, Elchalal U, Ginosar Y. Initial trauma management in advanced pregnancy. *Anesthesiology Clinics*. 2007; 25(1):117.
28. Katz VL. Perimortem cesarean delivery: its role in maternal mortality. *Seminars in Perinatology*. 2012; 36(1):68-72.
29. Jeejeebhoy FM, Zelop CM, Windrim R, Carvalho JCA, Dorian P, Morrison LJ. Management of cardiac arrest in pregnancy: a systematic review. *Resuscitation*. 2011; 82(7):801-809.
30. Oxford CM, Ludmir J. Trauma in pregnancy. *Clinical Obstetrics & Gynecology*. 2009; 52(4):611-629.
31. Brun PM, Chenaitia H, Dejesus I, Bessereau J, Bonello L, Pierre B. Ultrasound to perimortem caesarean delivery in prehospital settings. *Injury*. 2013; 44(1):151-152.
32. Criddle LM. Trauma in pregnancy: trauma care priorities don't change when the patient is pregnant. *American Journal of Nursing*. 2009; 109(11):41-48.
33. Ruffolo DC. Trauma care and managing the injured pregnant patient. *Journal of Obstetric, Gynecologic, and Neonatal Nursing* 2009; 38(6):704-714.
34. Queensland Clinical Guidelines. Postpartum haemorrhage. Guideline No. MN12.1-V4-R17. Queensland Health. 2012. Available from: <http://www.health.qld.gov.au/qcgl/>.
35. Queensland Clinical Guidelines. Intrapartum fetal surveillance. Guideline No. MN10.15-V3-R15. Queensland Health. 2010. Available from: <http://www.health.qld.gov.au/qcgl/>.
36. Afors K, Chandharan E. Use of continuous electronic fetal monitoring in a preterm fetus: clinical dilemmas and recommendations for practice. *Journal of Pregnancy*. 2011:1-7.
37. McGoldrick NP, Green C, Burke N, Quinlan C, McCormack D. Pregnancy and the orthopaedic patient. *Orthopaedics and Trauma*. 2012; 26(3):212-219.
38. Sadro C, Bernstein MP, Kanal KM. Imaging of trauma: Part 2, Abdominal trauma and pregnancy-a radiologist's guide to doing what is best for the mother and baby. *Am J Roentgenol*. 2012; 199(6):1207-19.
39. National Council on Radiation Protection and Measurements. Preconception and prenatal radiation exposure: Health effects and protective guidance, Report No.174. 2013.
40. Tremblay E, Thérasse E, Thomassin-Naggara I, Trop I. Quality initiatives: guidelines for use of medical imaging during pregnancy and lactation. *Radiographics*. 2012; 32(3):897-911.
41. Australian Radiation Protection and Nuclear Safety Agency. Radiation protection in diagnostic and interventional radiology; Radiation protection series RPS 14.1. 2008.
42. Australian Radiation Protection and Nuclear Safety Agency. Code of practice for radiation protection in the medical applications of ionizing radiation; Radiation protection series RPS 14. 2008.
43. Queensland Government. Risks to the fetus from diagnostic x-rays factsheet. Document number 20418-V5.0. Quality Information System. 2013 [cited 2013 November 21]. Available from: <http://qis.health.qld.gov.au/DocumentManagement/Default.aspx?DocumentID=20418&DocumentInstanceId=78584>.

44. Royal College of Obstetricians and Gynaecologists. Antepartum haemorrhage. Green-top Guideline No. 63. 2011.
45. Solomon N, Playforth K, Reynolds EW. Fetal-maternal hemorrhage: a case and literature review. *AJP Rep.* 2012; 2(1):7-14.
46. Wylie BJ, D'Alton ME. Fetomaternal hemorrhage. *Obstet Gynecol.* 2010; 115(5):1039-51.
47. Pearlman MD, Tintinalli JE, Lorenz RP. A prospective controlled study of outcome after trauma during pregnancy. *Am J Obstet Gynecol.* 1990; 162(6):1502-10.
48. Australian & New Zealand Society of Blood Transfusion Inc. Guidelines for laboratory assessment of fetomaternal haemorrhage. 2002 [cited 2013 July 01]. Available from: <http://www.anzsbts.org.au/publications/index.cfm>
49. Muench MV, Baschat AA, Reddy UM, Mighty HE, Weiner CP, Scalea TM, et al. Kleihauer-betke testing is important in all cases of maternal trauma. *J Trauma.* 2004; 57(5):1094-8.
50. Meguerdichian D. Complications in late pregnancy. *Emergency Medicine Clinics of North America.* 2012; 30(4):919-936.
51. Dhanraj D, Lambers D. The incidences of positive Kleihauer-Betke test in low-risk pregnancies and maternal trauma patients. *Am J Obstet Gynecol.* 2004; 190(5):1461-3.
52. Goodwin TM, Breen MT. Pregnancy outcome and fetomaternal hemorrhage after noncatastrophic trauma. *Am J Obstet Gynecol.* 1990; 162(3):665-71.
53. Rose PG, Strohm PL, Zuspan FP. Fetomaternal hemorrhage following trauma. *Am J Obstet Gynecol.* 1985; 153(8):844-7.
54. Cabral AC, Reis ZS, Apocalypse IG, Osanan GC, Lage EM, Leite HV. Combined use of the cardiofemoral index and middle cerebral artery Doppler velocimetry for the prediction of fetal anemia. *Int J Gynaecol Obstet.* 2010; 111(3):205-8.
55. Schenone MH, Mari G. The MCA Doppler and its role in the evaluation of fetal anemia and fetal growth restriction. *Clin Perinatol.* 2011; 38(1):83-102.
56. National Blood Authority Australia. Guidelines on the prophylactic use of Rh D immunoglobulin (anti D) in Obstetrics. 2003 [cited 2013 July 01]. Available from: <http://www.blood.gov.au>.
57. The Royal Australian and New Zealand College of Obstetricians and Gynaecologists. Guidelines for the use of Rh (D) Immunoglobulin (Anti-D) in obstetrics in Australia. College Statement C-Obs 6. 2011.
58. Queensland Clinical Guidelines. Preterm labour. Guideline No. MN09.6-V4-R14. Queensland Health. 2009. Available from: <http://www.health.qld.gov.au/qcgl/>.
59. Schiff M, Holt V. Pregnancy outcomes following hospitalization for motor vehicle crashes in Washington State from 1989 to 2001. *American Journal Epidemiology.* 2005; 161(6):503.
60. Raja AS, Zabbo CP. Trauma in pregnancy. *Emergency Medicine Clinics of North America.* 2012; 30(4):937-948.
61. Brown HL. Trauma in pregnancy. *Obstet Gynecol.* 2009; 114(1):147-60.
62. National Blood Authority Australia. Patient blood management guidelines: module 1 - critical bleeding/massive transfusion 2011 [cited 2013 July 16]. Available from: <http://www.blood.gov.au>
63. Queensland Clinical Guidelines. Venous thromboembolism (VTE) prophylaxis in pregnancy and the puerperium. Guideline No. MN14.9-V4-R19. Queensland Health. 2014. Available from: <http://www.health.qld.gov.au/qcgl/>.

## Appendix A: Classification of major trauma in pregnancy

If any ONE criterion (except systolic BP\*) is present from any category (vital signs, injury pattern or mechanism of injury), consider the trauma 'Major' and respond accordingly.

<input checked="" type="checkbox"/> Vital signs criteria		
	Conscious state	Altered level of consciousness
	Respiratory rate	< 10 or > 30 breaths/minute
	SpO <sub>2</sub> (room air)	< 95%
	Heart rate	> 120 bpm
	*Systolic BP	< 90 mmHg
<i>*Interpret BP in conjunction with gestation, other vital signs, injury pattern and mechanism of injury</i>		
<input checked="" type="checkbox"/> Injury pattern criteria		
	Penetrating or blast injury to the head, neck, chest, abdomen, pelvis, axilla or groin	
	Significant blunt injury to a single region of head, neck, chest, abdomen, pelvis or axilla	
	Injury to any two or more body regions of head, neck, chest, abdomen, pelvis or axilla	
	Limb amputation above the wrist or ankle	
	Suspected spinal cord injuries	
	Burns > 20% or other complicated burn injury including burn injury to the hand, face, genitals, airway and respiratory tract	
	Serious crush injury	
	Major compound fracture or open dislocation with vascular compromise	
	Fractured pelvis	
	Fractures involving two or more of the following: femur, tibia, humerus	
<input checked="" type="checkbox"/> Mechanism of injury criteria		
	Ejected from vehicle	
	Fall from height > 3 metres	
	Involved in an explosion	
	Involved in a high impact motor vehicle crash with incursion into the occupants compartment	
	Involved in a vehicle rollover	
	Involved in a road traffic collision in which there was a fatality in the same vehicle	
	Entrapped for > 30 minutes	
	Pedestrian impact	
	Motorcyclist impact > 30 kph	

Adapted from: Queensland Government. Queensland Ambulance Service (QAS) Field Reference Guide. 2011 and Queensland Government, Statewide Clinical Coordination and Retrieval Services. SOP No.3.7 Criteria for early notification of trauma for interfaculty transfers



## **Appendix B: Perimortem caesarean section procedure**

*Large vertical abdominal incision required. Uterine incision may be either vertical or horizontal*




Image produced by: Herston Multimedia Unit, Metro North Hospital and Health Service, Queensland.

## Appendix C: Haemodynamic and laboratory values in pregnancy

### Mean values for haemodynamic changes throughout pregnancy

	Pre-pregnancy	1 <sup>st</sup> Trimester	2 <sup>nd</sup> Trimester	3 <sup>rd</sup> Trimester
Heart rate (beats/min)	70	78	82	85
Systolic BP (mmHg)	125	112	122	115
Diastolic BP (mmHg)	70	60	63	70
Central venous pressure (mmHg)	9.0	7.5	4.0	3.8
Femoral venous pressure (mmHg)	6	6	18	18
Cardiac output (L/min)	4.5	4.5	6.0	6.0
Uterine blood flow (mL/min)	4000	4200	5000	5600

Source: Suresh MS, Latoya Mason C, Munnur U. Cardiopulmonary resuscitation and the parturient. Best Practice and Research: Clinical Obstetrics and Gynaecology. 2010; 24(3):383-400.

### Pathology Queensland reference intervals

	Gestation (weeks)	Reference range	Units
White Blood Cells (WBC)	1–12	5.7–13.6	$\times 10^9/L$
	13–24	6.2–14.8	$\times 10^9/L$
	25–42	5.9–16.9	$\times 10^9/L$
	>42	5.7–16.9	$\times 10^9/L$
Neutrophils	1–12	3.6–10.1	$\times 10^9/L$
	13–24	3.8–12.3	$\times 10^9/L$
	25–42	3.9–13.1	$\times 10^9/L$
	>42	3.6–13.1	$\times 10^9/L$
Eosinophils	1–>42	<0.6	$\times 10^9/L$
Lymphocytes	1–12	1.1–3.5	$\times 10^9/L$
	13–24	0.9–3.9	$\times 10^9/L$
	25–42	1.0–3.6	$\times 10^9/L$
	>42	0.9–3.9	$\times 10^9/L$
Platelets	1–12	170–390	$\times 10^9/L$
	13–24	170–410	$\times 10^9/L$
	25–42	150–430	$\times 10^9/L$
	>42	150–430	$\times 10^9/L$
Red Blood Cells (RBC)	1–12	3.52–4.52	$\times 10^{12}/L$
	13–24	3.20–4.41	$\times 10^{12}/L$
	25–42	3.10–4.44	$\times 10^{12}/L$
	>42	3.10–4.52	$\times 10^{12}/L$
Haemoglobin	1–12	110–143	g/L
	13–24	100–137	g/L
	24–42	98–137	g/L
	>42	98–143	g/L
Haematocrit	1–12	0.31–0.41	
	13–24	0.30–0.38	
	25–42	0.28–0.39	
	>42	0.28–0.41	
Mean Cell Haemoglobin (MCH)	1–>42	27.5–33.0	pg
Mean Cell Haemoglobin Concentration (MCHC)	1–>42	320–360	g/L
Erythrocyte Sedimentation Rate (ESR)	1–12	<30	mm/hr
	13–24	<64	mm/hr
	>24	<72	mm/hr
Bicarbonate (Total CO <sub>2</sub> )	All	18–26	mmol/L
Creatinine	All	40–80	mmol/L
Protein (Total)	14–40	61–75	g/L
Albumin	27–40	33–40	g/L
Urate	1–14	0.10–0.25	mmol/L
	15–27	0.10–0.30	mmol/L
	>27	0.10–0.35	mmol/L

## Appendix D: Seat belt positioning in pregnancy

Correct positioning of the seat belt includes:

- Lap belt over hips below uterus
- Sash between breasts above uterus

Correct application of the seat belt

- Reduces maternal/fetal injuries
- Reduces ejection mortalities
- Improves fetal survival

Use of a lap belt only is not recommended. It increases uterine flexion and may increase placental abruption

*Correct and incorrect positioning of seat belt*




Image produced by: Herston Multimedia Unit, Metro North Hospital and Health Service, Queensland.

## Appendix E: Estimation of gestation

Measure the vertical distance in the midline from the symphysis pubis to the top of the fundus in centimetres. This measurement correlates approximately with the gestational age.

Considerations that may impact on accuracy include:

- Multiple pregnancy
- Growth restriction
- Poly/oligohydramnios
- Breech or abnormal lie

*Estimating gestational age by fundal height*







Image produced by: Herston Multimedia Unit, Metro North Hospital and Health Service, Queensland.

## Appendix F: Left lateral tilt positioning


*Inferior vena cava compression when positioned supine*



*Left lateral tilt (right side up) 15-30 degrees to relieve compression*



*Manual displacement of the uterus to relieve compression*



Images produced by: Herston Multimedia Unit, Metro North Hospital and Health Service, Queensland.

## Appendix G: Approximate fetal effective doses (mSv) from common radiological examinations

Examination	1 <sup>st</sup> Trimester	3 <sup>rd</sup> Trimester
<b>Conventional radiography</b>		
Skull	<0.01	<0.01
Chest	<0.01	<0.01
Cervical spine	<0.01	<0.01
Thoracic spine	<0.01	<0.01
Lumbar spine	2	6
Abdomen	1.5	2.5
Pelvis	1	2
Intravenous pyelogram (IVP)	2	10
Extremities	<0.01	<0.01
Mammography	<0.01	<0.01
Barium meal	1	6
Barium enema	7	25
<b>Computerised Tomography (CT)</b>		
Head	<0.005	<0.005
Neck	<0.005	<0.01
Chest without portal phase	0.1	0.6
Chest with portal phase	1	7
Chest (pulmonary embolism)	0.1	0.4
Chest/abdomen/pelvis	12	13
Abdomen/pelvis – single phase	12	12
Abdomen/pelvis – multiple phase	15	30
Thoracic spine	0.2	1.0
Lumbar spine	10	25
Pelvimetry	–	0.2

**Note: All doses should be treated as indicative only as individual doses can differ from the tabulated values by as much as a factor of 10, except for those examinations remote from the lower abdomen**

Source: Australian Radiation Protection and Nuclear Safety Agency. Radiation protection in diagnostic and interventional radiology; Radiation protection series RPS 14.1. 2008.

## **Acknowledgements**

Queensland Clinical Guidelines gratefully acknowledge the contribution of Queensland clinicians and other stakeholders who participated throughout the guideline development process particularly:

### **Working Party Co-clinical Leads**

Associate Professor Rebecca Kimble, Director Obstetric Services, Royal Brisbane and Women's Hospital

Associate Professor Daryl Wall, Director Trauma Services, Royal Brisbane and Women's Hospital

Dr Frances Williamson, Emergency Physician, Royal Brisbane and Women's Hospital

### **Working Party Members**

Ms Stephanie Azri, Clinical Social Worker, Metro South Hospital and Health Service

Ms Michelle Barrett, Clinical Nurse Consultant, Retrieval Services Queensland

Dr John Burke, Emergency Physician, Royal Brisbane and Women's Hospital

Ms Katie Burke, Trauma Care Coordinator, Trauma Service, Royal Brisbane and Women's Hospital

Ms Dale Daly-Watkins, Nursing Director, Trauma Service, Royal Brisbane and Women's Hospital

Ms Tegan Draheim, Medical Officer, Royal Brisbane and Women's Hospital

Professor Nick Fisk, Executive Dean, Faculty of Health Sciences, University of Queensland

Mr Michael Handy, Trauma Care Coordinator, Trauma Service, Royal Brisbane and Women's Hospital

Dr Catherine Hurn, Emergency Physician, Royal Brisbane and Women's Hospital

Dr Benjamin Keir, Radiation Safety Officer, Senior Medical Physicist, Biomedical Technology Services

Dr Duncan McAuley, Emergency Physician, Royal Brisbane and Women's Hospital

Dr Tom McHattie, Clinical Director, Obstetrics and Gynaecology, Bundaberg Base Hospital

Associate Professor, Cliff Pollard, Board Member, Metro North Hospital and Health Board

Dr Stephen Rashford, Medical Director, Queensland Ambulance Service

Ms Tish Ryder, Consumer Representative, Maternity Coalition Queensland

Ms Rhonda Taylor, Midwifery Unit Manager, The Townsville Hospital, Townsville

Dr Edward Weaver, Staff Specialist, Obstetrics and Gynaecology, Nambour Hospital

Dr Neil Widdicombe, Intensivist, Royal Brisbane and Women's Hospital

### **Queensland Clinical Guidelines Team**

Associate Professor Rebecca Kimble, Director

Ms Jacinta Lee, Program Manager

Ms Lyndel Gray, Clinical Nurse Consultant

Dr Brent Knack, Program Officer

Ms Jeanette Tyler, Clinical Nurse Consultant

### **Funding**

This clinical guideline was funded by Queensland Health, Health Systems Innovation Branch.